

- 2-1.42 Describe the indications, contraindications, advantages, disadvantages, complications, and technique for inserting an oropharyngeal and nasopharyngeal airway (C-1)
- 2-1.43 Describe the indications, contraindications, advantages, disadvantages, complications, and technique for ventilating a patient by: (C-1)
 - a. Mouth-to-mouth
 - b. Mouth-to-nose
 - c. Mouth-to-mask
 - d. One person bag-valve-mask
 - e. Two person bag-valve-mask
 - f. Three person bag-valve-mask
 - g. Flow-restricted, oxygen-powered ventilation device
- 2-1.44 Explain the advantage of the two person method when ventilating with the bag-valve-mask. (C-1)
- 2-1.45 Compare the ventilation techniques used for an adult patient to those used for pediatric patients. (C-3)
- 2-1.46 Describe indications, contraindications, advantages, disadvantages, complications, and technique for ventilating a patient with an automatic transport ventilator (ATV). (C-1)
- 2-1.47 Explain safety considerations of oxygen storage and delivery. (C-1)
- 2-1.48 Identify types of oxygen cylinders and pressure regulators (including a high-pressure regulator and a therapy regulator). (C-1)
- 2-1.49 List the steps for delivering oxygen from a cylinder and regulator. (C-1)
- 2-1.50 Describe the use, advantages and disadvantages of an oxygen humidifier. (C-1)
- 2-1.51 Describe the indications, contraindications, advantages, disadvantages, complications, liter flow range, and concentration of delivered oxygen for supplemental oxygen delivery devices. (C-3)
- 2-1.52 Define, identify and describe a tracheostomy, stoma, and tracheostomy tube. (C-1)
- 2-1.53 Define, identify, and describe a laryngectomy. (C-1)
- 2-1.54 Define how to ventilate with a patient with a stoma, including mouth-to-stoma and bag-valve-mask-to-stoma ventilation. (C-1)
- 2-1.55 Describe the special considerations in airway management and ventilation for patients with facial injuries. (C-1)
- 2-1.56 Describe the special considerations in airway management and ventilation for the pediatric patient. (C-1)
- 2-1.57 Differentiate endotracheal intubation from other methods of advanced airway management. (C-3)
- 2-1.58 Describe the indications, contraindications, advantages, disadvantages and complications of endotracheal intubation. (C-1)
- 2-1.59 Describe laryngoscopy for the removal of a foreign body airway obstruction. (C-1)
- 2-1.60 Describe the indications, contraindications, advantages, disadvantages, complications, equipment, and technique for direct laryngoscopy. (C-1)
- 2-1.61 Describe visual landmarks for direct laryngoscopy. (C-1)
- 2-1.62 Describe use of cricoid pressure during intubation. (C-1)
- 2-1.63 Describe indications, contraindications, advantages, disadvantages, complications, equipment and technique for digital endotracheal intubation. (C-1)
- 2-1.64 Describe the indications, contraindications, advantages, disadvantages, complications, equipment and technique for using a dual lumen airway. (C-3)
- 2-1.65 Describe the indications, contraindications, advantages, disadvantages, complications and equipment for rapid sequence intubation with neuromuscular blockade. (C-1)
- 2-1.66 Identify neuromuscular blocking drugs and other agents used in rapid sequence intubation. (C-1)
- 2-1.67 Describe the indications, contraindications, advantages, disadvantages, complications and equipment for sedation during intubation. (C-1)
- 2-1.68 Identify sedative agents used in airway management. (C-1)
- 2-1.69 Describe the indications, contraindications, advantages, disadvantages, complications, equipment and technique for nasotracheal intubation. (C-1)
- 2-1.70 Describe the indications, contraindications, advantages, disadvantages and complications for performing

United States Department of Transportation
National Highway Traffic Safety Administration
Paramedic: National Standard Curriculum

DECLARATIVE

1. The body's need for oxygen
2. Primary objective of emergency care
 - a. Ensure optimal ventilation
 - (1) Delivery of oxygen
 - (2) Elimination of CO₂
3. Brain death occurs within 6 to 10 minutes
4. Major prehospital causes of preventable death
 - a. Early detection
 - b. Early intervention
 - c. Lay-person BLS education
5. Most often neglected of prehospital skills
 - a. Basics taken for granted
 - b. Poor techniques
 - (1) BVM seal
 - (2) Improper positioning
 - (3) Failure to reassess

1. Function of the upper airway
 - a. Warm
 - b. Filter
 - c. Humidify
2. Pharynx
 - a. Nasopharynx
 - (1) Formed by the union of facial bones
 - (2) Orientation of nasal floor is towards the ear not the eye
 - (3) Separated by septum
 - (4) Lined with
 - (a) Mucous membranes
 - (b) Cilia
 - (5) Turbinate
 - (a) Parallel to nasal floor
 - (b) Provide increased surface area for air
 - i) Filtration
 - ii) Humidifying
 - iii) Warming
 - (6) Sinuses
 - (a) Cavities formed by cranial bones
 - (b) Appear to further trap bacteria and act as tributaries for fluid to and from Eustachian tubes and tear ducts
 - i) Commonly become infected
 - ii) Fracture of certain sinus bones may cause cerebrospinal fluid (CSF) leak
 - (7) Tissues extremely delicate and vascular
 - (a) Improper or overly aggressive placement of tubes or airways will cause significant bleeding which may not be controlled by direct pressure

United States Department of Transportation
National Highway Traffic Safety Administration
Paramedic: National Standard Curriculum

[illegible]

- b. Oropharynx
 - (1) Teeth
 - (a) 32 adult
 - (b) Requires significant force to dislodge
 - (c) May fracture or avulse causing obstruction
 - (2) Tongue
 - (a) Large muscle attached at the mandible and hyoid bones
 - (b) Most common airway obstruction
 - (3) Palate
 - (a) Roof of mouth separates oro/ nasopharynx
 - i) Anterior is hard palate
 - ii) Posterior (beyond the teeth) is soft palate
 - (4) Adenoids
 - (a) Lymph tissue located in the mouth and nose that filters bacteria
 - (b) Frequently infected and swollen
 - (5) Posterior tongue
 - (6) Epiglottis
 - (7) Vallecula
 - (a) "Pocket" formed by the base of the tongue and epiglottis
 - (b) Important landmark for endotracheal intubation
- 3. Larynx
 - a. Attached to hyoid bone
 - (1) "Horseshoe-shaped" bone between the chin and mandibular angle
 - (2) Supports trachea
 - (3) Made of cartilage
 - b. Thyroid cartilage
 - (1) First tracheal cartilage
 - (2) "Shield-shaped"
 - (a) Cartilage anterior
 - (b) Smooth muscle posterior
 - (3) Laryngeal prominence
 - (a) "Adam's Apple" anterior prominence of thyroid cartilage
 - (b) Glottic opening directly behind
 - c. Glottic opening
 - (1) Narrowest part of adult trachea
 - (2) Patency heavily dependent on muscle tone
 - (3) Contain vocal bands
 - (a) White bands of cartilage
 - (b) Produce voice
 - d. Arytenoid cartilage
 - (1) "Pyramid-like" posterior attachment of vocal bands
 - (2) Important landmark for endotracheal intubation
 - e. Pyriform fossae
 - (1) "Hollow pockets" along the lateral borders of the larynx
 - f. Cricoid ring
 - (1) First tracheal ring
 - (2) Completely cartilaginous
 - (3) Compression occludes esophagus (Sellick maneuver)
 - g. Cricothyroid membrane
 - (1) Fibrous membrane between cricoid and thyroid cartilage

))))))))))

b.	Oxygen	104.0 torr (13.7%)
c.	CO ₂	40.0 torr (5.2%)
d.	Water	47.0 torr (6.2%)

1. Definition - the number of times a person breathes in one minute
2. Neural control
 - a. Primary control from the medulla and pons
 - b. Medulla
 - (1) Primary involuntary respiratory center
 - (2) Connected to respiratory muscles by vagus nerve
 - c. Pons
 - (1) Apneustic center - secondary control center if medulla fails to initiate respiration
 - (2) Pneumotaxic center - controls expiration
3. Chemical stimuli
 - a. Receptors for O_2 / CO_2 balance
 - (1) Cerebrospinal fluid pH
 - (2) Carotid bodies (sinus)
 - (3) Aortic arch
 - b. Hypoxic drive - respiratory stimulus dependent on O_2 rather than CO_2 in the blood
4. Control of respiration by other factors
 - a. Body temperature - respirations increase with fever
 - b. Drug and medications - may increase or decrease respirations depending on their physiologic action
 - c. Pain - increases respirations
 - d. Emotion - increases respirations
 - e. Hypoxia - increases respirations
 - f. Acidosis - respirations increase as compensatory response to increased CO_2 production
 - g. Sleep - respirations decrease

1. Obstruction
 - a. Tongue
 - (1) Most common airway obstruction
 - (2) Snoring respirations
 - (3) Corrected with positioning
 - b. Foreign body
 - (1) May cause partial or full obstruction
 - (2) Symptoms include
 - (a) Choking
 - (b) Gagging
 - (c) Stridor
 - (d) Dyspnea
 - (e) Aphonia (unable to speak)
 - (f) Dysphonia (difficulty speaking)
 - c. Laryngeal spasm and edema
 - (1) Spasm
 - (a) Spasmodic closure of vocal cords
 - (b) Most frequently caused by
 - i) Trauma from over aggressive technique during intubation

United States Department of Transportation
National Highway Traffic Safety Administration

[illegible]

- United States Department of Transportation
National Highway Traffic Safety Administration
Paramedic: National Standard Curriculum

[illegible]

- United States Department of Transportation
National Highway Traffic Safety Administration
Paramedic: National Standard Curriculum

[illegible]

- United States Department of Transportation
National Highway Traffic Safety Administration
Paramedic: National Standard Curriculum

[illegible]

7. Flow-restricted, oxygen-powered ventilation devices
 - a. The valve opening pressure at the cardiac sphincter is approx 30 cm H₂O
 - b. These devices operate at or below 30 cm H₂O to prevent gastric distension
 - c. Indications
 - (1) Delivery of high volume/ high concentration of O₂ (1 L/ sec)
 - (2) Awake compliant patients
 - (3) Unconscious patient with caution
 - d. Contraindications
 - (1) Noncompliant patients
 - (2) Poor tidal volume
 - (3) Small children
 - e. Advantages
 - (1) Self administered
 - (2) Delivers high volume/ high concentration O₂
 - (3) O₂ delivered in response to inspiratory effort (no O₂ wasting)
 - (4) O₂ volume delivery is regulated by inspiratory effort minimizing overinflation risk
 - (5) O₂ volume delivery is also restricted to less than 30 cm H₂O
 - f. Disadvantages
 - (1) Cannot monitor lung compliance
 - (2) Requires O₂ source
 - g. Complications
 - (1) Gastric distension
 - (2) Barotrauma
 - h. Method
 - (1) Mask is held manually in place
 - (2) Negative pressure upon inspiration triggers O₂ delivery or medic triggers release button
 - (3) Patient is monitored for adequate tidal volume and oxygenation
8. Automatic transport ventilators
 - a. Volume/ rate controlled
 - b. Indications
 - (1) Extended ventilation of intubated patients
 - (2) In situations in which a BVM is used
 - (3) Can be used during CPR
 - c. Contraindications
 - (1) Awake patients
 - (2) Obstructed airway
 - (3) Increased airway resistance
 - (a) Pneumothorax (after needle decompression)
 - (b) Asthma
 - (c) Pulmonary edema
 - d. Advantages
 - (1) Frees personnel to perform other tasks
 - (2) Lightweight
 - (3) Portable
 - (4) Durable
 - (5) Mechanically simple
 - (6) Adjustable tidal volume
 - (7) Adjustable rate
 - (8) Adapts to portable O₂ tank

[illegible]

[illegible]

- e. Disadvantages
 - (1) Cannot detect tube displacement
 - (2) Does not detect increasing airway resistance
 - (3) Difficult to secure
 - (4) Dependent on O₂ tank pressure
- 9. Cricoid pressure - Sellick's maneuver
 - a. Pressure on cricoid Ring
 - b. Occludes esophagus
 - c. Facilitates intubation by moving the larynx posteriorly
 - d. Helps to prevent passive emesis
 - e. Can help minimize gastric distension during bag-valve-mask ventilation
 - f. Indications
 - (1) Vomiting is imminent or occurring
 - (2) Patient cannot protect own airway
 - g. Contraindications
 - (1) Use with caution in cervical spine injury
 - h. Advantages
 - (1) Noninvasive
 - (2) Protects from aspiration as long as pressure is maintained
 - i. Disadvantages
 - (1) May have extreme emesis if pressure is removed
 - (2) Second rescuer required for bag-valve-mask ventilation
 - (3) May further compromise injured cervical spine
 - j. Complications
 - (1) Laryngeal trauma with excessive force
 - (2) Esophageal rupture from unrelieved high gastric pressures
 - (3) Excessive pressure may obstruct the trachea in small children
 - k. Method
 - (1) Locate the anterior aspect of the cricoid ring
 - (2) Apply firm, posterior pressure
 - (3) Maintain pressure until the airway is secured with an endotracheal tube
- 10. Artificial ventilation of the pediatric patient
 - a. Flat nasal bridge makes achieving mask seal more difficult
 - b. Compressing mask against face to improve mask seal results in obstruction
 - c. Mask seal best achieved with jaw displacement (two person bag-valve-mask)
 - d. Bag-valve-mask ventilation
 - (1) Bag size
 - (a) Full-term neonates and infants - minimum of 450 ml tidal volume (pediatric BVM)
 - (b) Children up to eight years of age - pediatric BVM preferred but adult-sized BVM (1500 ml) may be used
 - (c) Children over eight years of age require adult-sized BVM for adequate ventilation
 - (d) Proper mask fit
 - (e) Length based resuscitation tape
 - (f) Bridge of nose to cleft of chin
 - (2) Proper mask position and seal (EC-clamp)
 - (a) Place mask over mouth and nose; avoid compressing the eyes
 - (b) Using one hand, place thumb on mask at apex and index finger on mask at chin (C-grip)

[illegible]

- a) May use larger tubes
 - b) May lavage more aggressively
 - c) Safe to pass in facial fracture
 - d) Avoids nasopharynx
- iv) Disadvantages
 - a) May interfere with visualization during Intubation
- v) Method
 - a) Neutral or flexed head position
 - b) Introduce tube down midline
 - c) Procedure same as NG
- vi) Complications
 - a) Same as NG
 - b) Patient may bite tube

1. Manual maneuvers

(1) Technique

- b. Jaw-thrust without head-tilt maneuver

- [illegible]

))))))

- United States Department of Transportation
National Highway Traffic Safety Administration
Paramedic: National Standard Curriculum

[illegible]

- United States Department of Transportation
National Highway Traffic Safety Administration
Paramedic: National Standard Curriculum

- xii) Confirm placement with multiple methods
- xiii) Reconfirm placement with major patient movement or head movement

- [illegible]

[illegible]

- United States Department of Transportation
National Highway Traffic Safety Administration
Paramedic: National Standard Curriculum

[illegible]

- United States Department of Transportation
National Highway Traffic Safety Administration
Paramedic: National Standard Curriculum

- (a) Rapid insertion
 - (b) No special equipment
 - (c) Does not require sniffing position
- (5) Disadvantages
 - (a) Impossible to suction trachea when tube is in esophagus
 - (b) Adults only
 - (c) Unconscious only
 - (d) Very difficult to intubate around
- (6) Method
 - (a) Head - neutral position
 - (b) Pre-intubation precautions
 - (c) Insert with jaw-lift at midline
 - (d) Inflate pharyngeal cuff with 100 ccs of air
 - (e) Inflate distal cuff with 10-15 ccs of air
 - (f) Ventilate through longest tube first (pharyngeal)
 - i) Chest rise indicates esophageal placement of distal tip
 - ii) No chest rise indicates tracheal placement, switch ports and ventilate
- (7) Special considerations
 - (a) Good assessment skills are essential to confirm proper placement
 - (b) Mis-identification of placement has been reported
 - (c) Reinforce multiple confirmation techniques

1. Sedation in emergency intubation
 - a. Sedatives are used in airway management to
 - (1) Reduce anxiety
 - (2) Induce amnesia
 - (3) Decrease the gag reflex
 - b. Indications
 - (1) Combative patients
 - (2) Patients who require aggressive airway management but who are too conscious to tolerate intubation
 - (3) Agitated patients
 - c. Contraindications
 - (1) Known sensitivity to the medications
 - d. Advantages
 - (1) Decreases anxiety
 - (2) Induces amnesia
 - e. Disadvantages
 - (1) Respiratory depression
 - (2) Vomiting/ aspiration
 - f. Pharmacology
 - (1) Decreases anxiety
 - (2) Increases patient compliance
 - (3) Often produces amnesia to procedure
 - (4) Enhances ease of intubation
 - (5) Types of medications used
 - (a) Haloperidol
 - (b) Barbiturates

United States Department of Transportation
National Highway Traffic Safety Administration

- (c) Benzodiazepines
- (d) Etomidate
- (e) Narcotics
- (f) Ketamine

- [illegible]

[illegible]

- (3) Tube may be removed/ cleaned and replaced
 - b. Stenosis
 - (1) Stoma spontaneously narrows
 - (a) Potentially life-threatening
 - (b) Soft tissue swelling decreases stoma diameter
 - (2) Trach tube is difficult or impossible to replace
 - (3) ET tube must be placed before total obstruction
 - c. Suctioning
 - (1) Must be done with extreme caution if laryngeal edema is suspected
 - (2) Procedure
 - (a) Preoxygenate
 - (b) Inject 3 cc sterile saline down trachea
 - (c) Instruct patient to exhale
 - (d) Insert suction catheter until resistance detected
 - (e) Instruct patient to cough or exhale
 - (f) Suction during withdrawal
 - d. Tube replacement
 - (1) Lubricate appropriately sized tracheostomy tube or ET tube (5.0 or larger)
 - (2) Instruct patient to exhale
 - (3) Gently insert tube about 1-2 cm beyond balloon cuff
 - (4) Inflate balloon cuff
 - (5) Confirm comfort, patency and proper placement
 - (6) Ensure false lumen was not created
- 2. Dental appliances
 - a. Dentures, partial plates, etc.
 - b. Best removed before intubation
- 3. Airway management considerations for patients with facial injuries
 - a. Facial injuries suggest the possibility of cervical spine injury
 - (1) In-line stabilization
 - (a) Trauma technique endotracheal intubation
 - b. Foreign body/ blood in oropharynx
 - (1) Suction airway
 - c. Inability to ventilate/ intubate orally
 - (1) Requires surgical intervention